

SUPERSYMMETRIC U(1) AT THE TEV SCALE

E. KEITH

*Center for Particle Physics, University of Texas
Austin, TX 78712, USA*

ERNEST MA

*Department of Physics, University of California
Riverside, CA 92521, USA*

If there exists an arbitrary supersymmetric U(1) gauge factor at the TeV scale, under which the two Higgs superfields $H_{1,2}$ of the standard model are nontrivial, and if there is also a singlet superfield S such that the $H_1 H_2 S$ term is allowed in the superpotential, then the structure of the two-doublet Higgs sector at the electroweak scale is more general than that of the MSSM (Minimal Supersymmetric Standard Model). Under further assumptions of grand unification and universal soft supersymmetry breaking terms, the scale of U(1) breaking is related to the parameter $\tan \beta \equiv v_2/v_1$.

1 Two Doublets and a Singlet

Assume an arbitrary supersymmetric U(1) gauge factor at the TeV scale. Call it $U(1)_X$. Then under $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_X$, consider two doublet and one singlet Higgs superfields transforming as follows:¹

$$H_1 \sim (1, 2, -1/2; -a), \quad (1)$$

$$H_2 \sim (1, 2, 1/2; -1 + a), \quad (2)$$

$$S \sim (1, 1, 0; 1). \quad (3)$$

The superpotential of this model is then given by

$$W = f H_1 H_2 S + \dots \quad (4)$$

This has the advantage that the $\mu H_1 H_2$ term in the MSSM (Minimal Supersymmetric Standard Model) is replaced by $\mu = f \langle S \rangle$, which is a possible solution of the so-called μ problem. (Since the $H_1 H_2$ term is allowed by supersymmetry in the MSSM, there is no natural understanding as to why μ should not be very much larger than the scale of supersymmetry breaking. On the other hand, with $\langle S \rangle$ at the TeV scale and f a typical Yukawa coupling, it is natural for μ to be at the electroweak scale, leading to the possibility that all physical members of the two Higgs doublets are at the 100 GeV scale.)

In the scalar sector, let H_1 be represented by $\tilde{\Phi}_1 = (\bar{\phi}_1^0, -\phi_1^-)$ and H_2 by $\Phi_2 = (\phi_2^+, \phi_2^0)$, and S by χ^0 , then $\langle \chi^0 \rangle = u$ breaks only $U(1)_X$, and $\langle \phi_{1,2}^0 \rangle = v_{1,2}$

breaks $SU(2)_L \times U(1)_Y$ to $U(1)_Q$. The contribution of the fH_1H_2S term in the superpotential to the Higgs potential is given by

$$V_F = f^2[(\Phi_1^\dagger \Phi_2)(\Phi_2^\dagger \Phi_1) + (\Phi_1^\dagger \Phi_1 + \Phi_2^\dagger \Phi_2)\bar{\chi}\chi], \quad (5)$$

and the gauge contribution is

$$\begin{aligned} V_D &= \frac{1}{8}g_2^2[(\Phi_1^\dagger \Phi_1)^2 + (\Phi_2^\dagger \Phi_2)^2 + 2(\Phi_1^\dagger \Phi_1)(\Phi_2^\dagger \Phi_2) - 4(\Phi_1^\dagger \Phi_2)(\Phi_2^\dagger \Phi_1)] \\ &+ \frac{1}{8}g_1^2[\Phi_1^\dagger \Phi_1 - \Phi_2^\dagger \Phi_2]^2 \\ &+ \frac{1}{2}g_x^2[-a\Phi_1^\dagger \Phi_1 - (1-a)\Phi_2^\dagger \Phi_2 + \bar{\chi}\chi]^2. \end{aligned} \quad (6)$$

The supersymmetry breaking terms are contained in

$$V_{soft} = \mu_1^2 \Phi_1^\dagger \Phi_1 + \mu_2^2 \Phi_2^\dagger \Phi_2 + m^2 \bar{\chi}\chi + (fA_f \Phi_1^\dagger \Phi_2 \chi + h.c.). \quad (7)$$

First, we assume $u \sim M_{SUSY} \sim \text{TeV}$. This is natural because $U(1)_X$ cannot be broken without also breaking the supersymmetry. Of course, it is also possible² to have $u < \text{TeV}$. Second, we assume that at the 100 GeV scale, there are just the two Higgs doublets. This requires $fA_f u = m_{12}^2 \ll (\text{TeV})^2$.

2 Reduced Higgs Potential

With $\langle \chi \rangle = u$, the scalar field $\sqrt{2}Re\chi$ is physical and has a mass given by $m^2 = 2g_x^2 u^2$. The cubic $(\Phi_1^\dagger \Phi_1)\sqrt{2}Re\chi$ coupling is then $\sqrt{2}u(f^2 - g_x^2 a)$. Hence the effective quartic $(\Phi_1^\dagger \Phi_1)^2$ coupling is given by

$$\begin{aligned} \lambda_1 &= \frac{1}{4}(g_1^2 + g_2^2) + g_x^2 a^2 - \frac{2u^2(f^2 - g_x^2 a)^2}{2g_x^2 u^2} \\ &= \frac{1}{4}(g_1^2 + g_2^2) + 2af^2 - \frac{f^4}{g_x^2}. \end{aligned} \quad (8)$$

Similarly, the other effective quartic scalar couplings are as follows:

$$\lambda_2 = \frac{1}{4}(g_1^2 + g_2^2) + 2(1-a)f^2 - \frac{f^4}{g_x^2}, \quad (9)$$

$$\lambda_3 = -\frac{1}{4}g_1^2 + \frac{1}{4}g_2^2 + f^2 - \frac{f^4}{g_x^2}, \quad (10)$$

$$\lambda_4 = -\frac{1}{2}g_2^2 + f^2. \quad (11)$$

Note that in the limit $f = 0$, we recover the Higgs structure of the MSSM.

Let $\tan \beta \equiv v_2/v_1$ and $v \equiv (v_1^2 + v_2^2)^{1/2}$, then

$$(m_h^2)_{max} = 2v^2[\lambda_1 \cos^4 \beta + \lambda_2 \sin^4 \beta + 2(\lambda_3 + \lambda_4) \sin^2 \beta \cos^2 \beta] + \epsilon, \quad (12)$$

where the radiative correction due to the t quark and squarks is given by

$$\epsilon \simeq \frac{3g_s^2 m_t^4}{8\pi^2 M_W^2} \ln \left(1 + \frac{\tilde{m}^2}{m_t^2} \right). \quad (13)$$

Hence the existence of an extra supersymmetric U(1) gauge factor at the TeV scale implies

$$(m_h^2)_{max} = M_Z^2 \cos^2 2\beta + \epsilon + \frac{f^2}{\sqrt{2}G_F} \left[A - \frac{f^2}{g_x^2} \right], \quad (14)$$

where

$$A = \frac{3}{2} + (2a - 1) \cos 2\beta - \frac{1}{2} \cos^2 2\beta. \quad (15)$$

If $A > 0$, the MSSM bound can be exceeded. However, for a given g_x^2 , f^2 is still bounded from the requirement that the reduced Higgs potential for the two doublets be bounded from below. Hence $(m_h^2)_{max}$ is bounded. See Figure 1 of Ref. 1. For specific models derivable from E_6 , the upper bound is raised to no higher than about 146 GeV as compared to 128 GeV in the MSSM. For $g_x^2 = 0.5$, the absolute upper bound is about 190 GeV.

3 Models Based on E_6

Consider the sequential reduction of E_6 :

$$E_6 \rightarrow SO(10) [\times U(1)_\psi], \quad (16)$$

$$SO(10) \rightarrow SU(5) [\times U(1)_\chi], \quad (17)$$

$$SU(5) \rightarrow SU(3)_C \times SU(2)_L [\times U(1)_Y]. \quad (18)$$

Assuming that a single extra U(1) survives down to the TeV energy scale, it is generally given by a linear combination of $U(1)_\psi$ and $U(1)_\chi$ which we call $U(1)_\alpha$. Under the maximal subgroup $SU(3)_C \times SU(3)_L \times SU(3)_R$, the fundamental representation of E_6 is given by

$$\mathbf{27} = (3, 3, 1) + (3^*, 1, 3^*) + (1, 3^*, 3). \quad (19)$$

Under the subgroup $SU(5) \times U(1)_\psi \times U(1)_\chi$, we then have

$$\begin{aligned} \mathbf{27} = & (10; 1, -1)[(u, d), u^c, e^c] + (5^*; 1, 3)[d^c, (\nu_e, e)] + (1; 1, -5)[N] \\ & + (5; -2, 2)[h, (E^c, N_E^c)] + (5^*; -2, -2)[h^c, (\nu_E, E)] + (1; 4, 0)[S], \end{aligned} \quad (20)$$

where the $U(1)$ charges refer to $2\sqrt{6}Q_\psi$ and $2\sqrt{10}Q_\chi$. Note that the known quarks and leptons are contained in $(10; 1, -1)$ and $(5^*; 1, 3)$, and the two Higgs scalar doublets are represented by (ν_E, E) and (E^c, N_E^c) . Let

$$Q_\alpha = Q_\psi \cos \alpha - Q_\chi \sin \alpha, \quad (21)$$

then the η -model³ is obtained with $\tan \alpha = \sqrt{3/5}$ and we have

$$\mathbf{27} = (10; 2) + (5^*; -1) + (1; 5) + (5; -4) + (5^*; -1) + (1; 5), \quad (22)$$

where $2\sqrt{15}Q_\eta$ is denoted. Hence $a = 1/5$ and $g_x^2 = (25/36)g_1^2$ for this model. If we take $\tan \alpha = -1/\sqrt{15}$, we get the N -model⁴ with

$$\mathbf{27} = (10; 1) + (5^*; 2) + (1; 0) + (5; -2) + (5^*; -3) + (1; 5), \quad (23)$$

where $2\sqrt{10}Q_N$ is denoted. Here $a = 3/5$ and $g_x^2 = (25/24)g_1^2$. As a last example, in the exotic left-right model⁵, $a = \tan^2 \theta_W$ and

$$g_x^2 = \frac{(g_1^2 + g_2^2)(1 - \sin^2 \theta_W)^2}{4(1 - 2 \sin^2 \theta_W)}. \quad (24)$$

4 Supersymmetric Scalar Masses

As a reasonable and predictive procedure, we will adopt the common hypothesis that soft supersymmetry-breaking operators appear at the grand-unification scale as the result of a hidden sector which is linked to the observable sector only through gravity. Hence these terms will be assumed to be universal, *i.e.* of the same magnitude for all fields. Consider now the masses of the supersymmetric scalar partners of the quarks and leptons:

$$m_B^2 = m_0^2 + m_R^2 + m_F^2 + m_D^2, \quad (25)$$

where m_0 is a universal soft supersymmetry-breaking mass, m_R^2 is a correction generated by the renormalization-group equations running from the grand-unification scale down to the TeV scale, m_F is the explicit mass of the fermion partner, and m_D^2 is a term induced by gauge symmetry breaking with rank reduction and can be expressed in terms of the gauge-boson masses. In the MSSM, m_D^2 is of order M_Z^2 and does not change m_B significantly. In the $U(1)_\alpha$ -extended model, m_D^2 is of order $M_{Z'}^2$, and will affect m_B in a nontrivial way. The contributions to m_D^2 from $U(1)_\alpha$ are

$$\Delta m_D^2(10; 1, -1) = \frac{1}{8}M_{Z'}^2 \left(1 + \sqrt{\frac{3}{5}} \tan \alpha \right), \quad (26)$$

$$\Delta m_D^2(5^*; 1, 3) = \frac{1}{8} M_{Z'}^2 \left(1 - 3\sqrt{\frac{3}{5}} \tan \alpha \right), \quad (27)$$

$$\Delta m_D^2(1; 1, -5) = \frac{1}{8} M_{Z'}^2 \left(1 + \sqrt{15} \tan \alpha \right), \quad (28)$$

$$\Delta M_D^2(5; -2, 2) = -\frac{1}{4} M_{Z'}^2 \left(1 + \sqrt{\frac{3}{5}} \tan \alpha \right), \quad (29)$$

$$\Delta m_D^2(5^*; -2, -2) = -\frac{1}{4} M_{Z'}^2 \left(1 - \sqrt{\frac{3}{5}} \tan \alpha \right), \quad (30)$$

$$\Delta m_D^2(1; 4, 0) = \frac{1}{2} M_{Z'}^2. \quad (31)$$

Thus it is actually possible⁶ for exotic scalar quarks and leptons to be lighter than the ordinary ones. Furthermore, since these masses are also present in the Higgs potential, the contributions of Δm_D^2 due to $U(1)_\alpha$ are essential in constraining its parameters, *i.e.* m_A and $\tan \beta$.

5 Matching of Parameters at the TeV Scale

In the two-doublet Higgs potential, the soft terms are given by

$$V_{soft} = m_1^2 \Phi_1^\dagger \Phi_1 + m_2^2 \Phi_2^\dagger \Phi_2 + m_{12}^2 (\Phi_1^\dagger \Phi_2 + \Phi_2^\dagger \Phi_1). \quad (32)$$

In the $U(1)_\alpha$ -extended model, they are related to M_Z and the pseudoscalar mass m_A as follows:

$$m_{12}^2 = -m_A^2 \sin \beta \cos \beta, \quad (33)$$

$$m_1^2 = m_A^2 \sin^2 \beta - \frac{1}{2} M_Z^2 \cos 2\beta - \frac{2f^2}{g_Z^2} M_Z^2 \left[2 \sin^2 \beta + \left(1 - \sqrt{\frac{3}{5}} \tan \alpha \right) \cos^2 \beta - \frac{3f^2}{2 \cos^2 \alpha g_\alpha^2} \right], \quad (34)$$

$$m_2^2 = m_A^2 \cos^2 \beta + \frac{1}{2} M_Z^2 \cos 2\beta - \frac{2f^2}{g_Z^2} M_Z^2 \left[2 \cos^2 \beta + \left(1 + \sqrt{\frac{3}{5}} \tan \alpha \right) \sin^2 \beta - \frac{3f^2}{2 \cos^2 \alpha g_\alpha^2} \right]. \quad (35)$$

In the limit $f = 0$ in the above, we recover the well-known results of the MSSM.

On the other hand, Eqs. (7) and (25) tell us that

$$m_{12}^2 = f A_f u, \quad (36)$$

$$m_1^2 = m_0^2 + m_{R1}^2 + f^2 u^2 - \frac{1}{4} \left(1 - \sqrt{\frac{3}{5}} \tan \alpha \right) M_{Z'}^2, \quad (37)$$

$$m_2^2 = m_0^2 + m_{R2}^2 + f^2 u^2 - \frac{1}{4} \left(1 + \sqrt{\frac{3}{5}} \tan \alpha \right) M_{Z'}^2, \quad (38)$$

where $M_{Z'}^2 = (4/3) \cos^2 \alpha g_\alpha^2 u^2$ and m_{R2}^2 differs from m_{R1}^2 in that the former contains the contribution from the t Yukawa coupling and the latter does not. Given a particular $U(1)_\alpha$ from E_6 , we can start at the grand-unification scale with m_0 , A_0 , and $m_{1/2}$, then for a given value of f , the matching of Eqs. (36) to (38) with Eqs. (33) to (35) will allow us to derive u and $\tan \beta$.

In our approach¹, we assume that the term $f' h h^c S$ in the superpotential is important enough to drive m_χ^2 in Eq. (7) negative, so that $U(1)_\alpha$ is broken with

$$M_{Z'} = \sqrt{-2m_\chi^2} = \frac{2}{\sqrt{3}} g_\alpha \cos \alpha |u|, \quad (39)$$

where g_α is assumed equal approximately to $\sqrt{5/3} g_1$. The mass of the exotic quark h is then given by $f'|u|$. As mentioned already at the end of Sec. 1, we assume also that $f A_f$ to be small compared to u . In a different approach², f' is assumed zero, but $f A_f$ is taken to be rather large. In that case, $v_{1,2}$ and u cannot be separated into two different scales. Also, there is no solution with universal soft supersymmetry-breaking terms.

In Figures 4 to 9 of Ref. 1, we show typical solutions of $\tan \beta$ and $|u|$ for various inputs of m_0 , A_0 , the gluino mass, and f . We find $M_{Z'}$ to be around 1 TeV, m_h around 2 TeV, and $\tan \beta$ around 4. These solutions are much more constrained than the ones in the MSSM because the free parameter $m_{12}^2 = B\mu$ is now replaced by $f A_f u$.

6 Conclusion

An extra supersymmetric $U(1)_\alpha$ gauge factor from E_6 is a good possibility at the TeV energy scale. Its existence implies that the two-doublet Higgs structure at around 100 GeV will be observably different from that of the MSSM. Supersymmetric scalar masses are also very different because of the large Δm_D^2 contributions. The $U(1)_\alpha$ breaking scale and the well-known parameter $\tan \beta \equiv v_2/v_1$ are closely related.

Acknowledgments

The presenter of this talk (E.M.) thanks Goran Senjanovic and Alexei Smirnov for the organization of this stimulating conference. This work was supported in

part by the U. S. Department of Energy under Grant No. DE-FG03-94ER40837.

References

1. E. Keith and E. Ma, Phys. Rev. **D56**, 7155 (1997).
2. M. Cvetič, D. A. Demir, J. R. Espinosa, L. Everett, and P. Langacker, Phys. Rev. **D56**, 2861 (1997).
3. K. S. Babu, C. Kolda, and J. March-Russel, Phys. Rev. **D54**, 4635 (1996).
4. E. Ma, Phys. Lett. **B380**, 286 (1996).
5. E. Ma, Phys. Rev. **D36**, 274 (1987); E. Ma and D. Ng, *ibid.* **D49**, 6164 (1994).
6. E. Keith, E. Ma, and B. Mukhopadhyaya, Phys. Rev. **D55**, 3111 (1997).